

Development of Bismuth and Iron Catalysts for Asymmetric Synthesis

Giovedì, 16 Gennaio, 2020

15:00

Aula II

Dipartimento di Chimica "Giacomo Ciamician"
Via F. Selmi 2
40126 – Bologna



Thierry Ollevier

Département de chimie
Université Laval, Québec, Canada



Development of Bismuth and Iron Catalysts for Asymmetric Synthesis

Thierry Ollevier

Université Laval, Québec, Canada

Various Lewis acids have been developed as green catalysts for asymmetric synthesis. Chiral metal complexes derived from iron, copper, and bismuth salts have been employed in selected asymmetric C–C, C–Si, C–N, and C–S bond-forming reactions, such as the Mukaiyama aldol, epoxide opening, Michael and Diels-Alder reactions.¹ Enantioselective oxidation catalysts will also be presented.² As part of our ongoing interest in the development of greener reaction conditions, we report alternate reaction solvent systems. These results will contribute to the development of green acid catalysis for asymmetric synthesis.

1. (a) Li, M.; Carreras, V.; Jalba, A.; Ollevier, T. *Org. Lett.* **2018**, *20*, 995–998. (b) Lauzon, S.; Keipour, H.; Gandon, V.; Ollevier, T. *Org. Lett.* **2017**, *19*, 6324–6327. (c) Ollevier, T.; Plancq., B. *Chem. Commun.* **2012**, *48*, 2289–2291.
2. Jalba, A.; Régnier, N.; Ollevier, T. *Eur. J. Org. Chem.* **2017**, 1628–1637.